arXiv:1208.0962v1 [math.OC] 4 Aug 2012

Complete Real Time Solution of the General Nonlinear Filtering Problem without Memory

Xue Luo and Stephen S.-T. Yau, IEEE Fellow

Abstract

It is well known that the nonlinear filtering problem has important applications in both military and commercial industries. The central problem of nonlinear filtering is to solve the Duncan-Mortensen-Zakai (DMZ) equation in real time and in a memoryless manner. In this paper, we will show that under very mild conditions (which essentially say that the growth of the observation is faster than the growth of the drift), the unique nonnegative weak solution of the most general DMZ equation can be approximated by the solution to that restricted on a large ball with Dirichlet boundary condition. The error of this approximation tends to zero exponentially as the radius of the ball tends to infinity. Moreover, the numerical solution on the large ball can be efficiently obtained by Yau-Yau's algorithm, which depends only on solving the observation-independent Kolmogorov forward equation on each time step. Equally important, we show that the numerical solution obtained by Yau-Yau's algorithm converges to the solution restricted on the large ball in the L^1 sense, and a precise convergence estimate is given. Numerical simulation to 1D cubic sensor is illustrated at the end.

I. INTRODUCTION

The Kalman filters have been widely used in many areas such as navigational and guidance systems, radar tracking, solar mapping, and satellite orbit determination. However, they are limited to where the linearity assumptions of the drift term and observation term are satisfied, as well as that the Gaussian assumption of initial value is required. In practice, the nonlinear drift/observation terms enter to the modeling naturally, say the cubic sensor. Therefore, it is

X. Luo is with the department of Mathematics,Statistics and Computer Science, University of Illinois at Chicago, Science and Engineering Offices (M/C 249), 851 S. Morgan Street, Chicago, IL 60607-7045 `xluo6@uic.edu`

S. S.-T. Yau is the faculty of Department of mathematical sciences, Tsinghua University, Beijing, 100084, P.R.China. He is the Emeritus Distinguished Professor of University of Illinois at Chicago `yau@uic.edu`

necessary and interesting to solve the nonlinear filtering problems. By “solving” a nonlinear filtering problem, it is to determine the conditional density $\rho(x,t)$ of the stochastic process x_t, named the state, given the observation history $\{y_s\colon 0 \leqslant s \leqslant t\}$, another stochastic process related to x_t. In the late 1960s, Duncan [2], Mortensen [5] and Zakai [10] independently derived the Duncan–Mortensen–Zakai (DMZ) equation, which the conditional probability density $\rho(x,t)$ satisfy. Henceforth, the central problem in nonlinear filtering theory is to solve the DMZ equation in real time and memoryless way.

In this paper, we consider the nonlinear filtering problem based on the signal observation model

$$\begin{cases} dx_t = f(x_t,t)dt + G(x_t,t)dv_t, \\ dy_t = h(x_t,t)dt + dw_t, \end{cases} \tag{1.1}$$

where x_t and f are n-vectors, G is an $n \times r$ matrix, and v_t is an r-vector Brownian motion process with $E[dv_t dv_t^T] = Q(t)dt$, y_t and h are m-vectors and w_t is an m-vector Brownian motion process with $E[dw_t dw_t^T] = S(t)dt$ and $S(t) > 0$. We refer to x_t as the state of the system at time t with x_0 satisfying some distribution and y_t as the observation at time t with $y_0 = 0$. We assume that $\{v_t, t \geq 0\}$, $\{w_t, t \geq 0\}$ and x_0 are independent.

The unnormalized density function $\sigma(x,t)$ of x_t conditioned on the observation history $Y_t = \{y_s : 0 \leq s \leq t\}$, at least formally, satisfies the DMZ equation

$$\begin{cases} d\sigma(x,t) = L\sigma(x,t)dt + \sigma(x,t)h^T(x,t)S^{-1}(t)dy_t \\ \sigma(x,0) = \sigma_0(x), \end{cases} \tag{1.2}$$

in which $\sigma_0(x)$ is the probability density of the initial state x_0, and

$$L(*) \equiv \frac{1}{2}\sum_{i,j=1}^{n} \frac{\partial^2}{\partial x_i \partial x_j}\left[\left(GQG^T\right)_{ij} *\right] - \sum_{i=1}^{n} \frac{\partial (f_i *)}{\partial x_i}. \tag{1.3}$$

The DMZ equation (1.2) is a stochastic partial differential equation due to the term dy_t. There is no easy way to derive a recursive algorithm for solving this equation. In real applications, one may be more interested in constructing robust state estimators from observed sample paths with some property of robustness. For each “given” observation, making an invertible exponential transformation [7]

$$\sigma(x,t) = \exp\left[h^T(x,t)S^{-1}(t)y_t\right]\rho(x,t), \tag{1.4}$$

the DMZ equation is transformed into a deterministic partial differential equation with stochastic coefficients, which we will refer as the "pathwise-robust" DMZ equation

$$\begin{cases} \dfrac{\partial\rho}{\partial t}(x,t)+\dfrac{\partial}{\partial t}(h^TS^{-1})^Ty_t\rho(x,t) \\ \qquad\qquad = \exp\left(-h^TS^{-1}y_t\right)\left[L-\dfrac{1}{2}h^TS^{-1}h\right]\left[\exp\left(h^TS^{-1}y_t\right)\rho(x,t)\right] \\ \rho(x,0)=\sigma_0(x). \end{cases} \tag{1.5}$$

Or equivalently,

$$\begin{cases} \dfrac{\partial\rho}{\partial t}(x,t)=\dfrac{1}{2}D_w^2\rho(x,t)+F(x,t)\cdot\nabla\rho(x,t)+J(x,t)\rho(x,t) \\ \rho(x,0)=\sigma_0(x), \end{cases} \tag{1.6}$$

where

$$D_w^2=\sum_{i,j=1}^n(GQG^T)_{ij}\frac{\partial^2}{\partial x_i\partial x_j}, \tag{1.7}$$

$$F(x,t)=\left[\sum_{j=1}^n\frac{\partial}{\partial x_j}\left(GQG^T\right)_{ij}+\sum_{j=1}^n(GQG^T)_{ij}\frac{\partial K}{\partial x_j}-f_i\right]_{i=1}^n, \tag{1.8}$$

$$\begin{aligned} J(x,t)=&-\frac{\partial}{\partial t}(h^TS^{-1})^Ty_t+\frac{1}{2}\sum_{i,j=1}^n\frac{\partial^2}{\partial x_i\partial x_j}\left(GQG^T\right)_{ij}+\sum_{i,j=1}^n\frac{\partial}{\partial x_i}\left(GQG^T\right)_{ij}\frac{\partial K}{\partial x_j} \\ &+\frac{1}{2}\sum_{i,j=1}^n(GQG^T)_{ij}\left[\frac{\partial^2K}{\partial x_i\partial x_j}+\frac{\partial K}{\partial x_i}\frac{\partial K}{\partial x_j}\right]-\sum_{i=1}^n\frac{\partial f_i}{\partial x_i}-\sum_{i=1}^nf_i\frac{\partial K}{\partial x_i}-\frac{1}{2}(h^TS^{-1}h), \end{aligned} \tag{1.9}$$

where

$$K(x,t)=h^T(x,t)S^{-1}(t)y_t. \tag{1.10}$$

The exact solution to (1.5) or (1.6), generally speaking, doesn't have a closed form. Hence, many mathematicians pay their effort on seeking an efficient algorithm to construct a good approximate solution. One of the direction is the splitting-up method to the DMZ equation, which is first introduced by Besoussan, Glowinski, and Rascanu [1]. Their method is limited to bounded drift term and observation term. Yau and Yau [9] developed a novel algorithm to the robust DMZ equation, which get rid of the boundedness of drift term and observation term to some mild growth conditions. They proved that the unique nonnegative solution of the robust DMZ equation can be approximated by that restricted on a large ball with Dirichlet boundary condition. However, they only show their algorithm works well where the drift/observation terms don't have explicit dependence on time t and the noises are all white.

In this paper, we will extend Yau-Yau's algorithm to the most general setting of the robust DMZ equation. We will not only give the theoretic error estimates, but also illustrate the numerical simulation to support our theoretic results. The difficulties are two folds: on one hand, the existence and uniqueness of the robust DMZ equation in the most general setting hasn't been established yet. The well-posedness of the robust DMZ equation, under some conditions, has been investigated by [6], [3], [9], etc.; on the other hand, the most general setting introduced more terms which leads to much more involved computations and analysis.

Let $\mathcal{P}_k = \{0 = \tau_0 \le \tau_1 \le \cdots \le \tau_k = T\}$ be a partition of $[0, T]$. Let ρ_i be a solution of the robust DMZ equation with y_t freezed at $t = \tau_{i-1}$, for $\tau_{i-1} \le t \le \tau_i$, $i = 1, 2, \cdots, k$

$$\begin{cases} \dfrac{\partial \rho_i}{\partial t}(x,t) + \dfrac{\partial}{\partial t}\left(h^T S^{-1}\right)^T y_{\tau_{i-1}} \rho_i(x,t) \\ \qquad\qquad = \exp\left(-h^T S^{-1} y_{\tau_{i-1}}\right)\left[L - \dfrac{1}{2} h^T S^{-1} h\right]\left[\exp\left(h^T S^{-1} y_{\tau_{i-1}}\right)\rho_i(x,t)\right] \\ \rho_1(x,0) = \sigma_0(x), \\ \text{or} \\ \rho_i(x,\tau_{i-1}) = \rho_{i-1}(x,\tau_{i-1}), \quad \text{for } i = 2, 3, \cdots, k. \end{cases} \tag{1.11}$$

Define the norm of the partition $\mathcal{P}_k$ by $|\mathcal{P}_k| = \sup_{1 \le i \le k}(\tau_i - \tau_{i-1})$. Intuitively, as $|\mathcal{P}_k| \to 0$, we have

$$\sum_{i=1}^{k} \chi_{[\tau_{i-1},\tau_i]}(t)\rho_i(x,t) \to \rho(x,t)$$

in some sense, for all $0 \le t \le T$, where $\rho(x,t)$ is the solution of the robust DMZ equation (1.5) or (1.6).

The proposition below helps to shift the computations off-line. This is the key ingredient of Yau-Yau's algorithm.

Proposition 1.1: For each $\tau_{i-1} \le t \le \tau_i$, $i = 1, 2, \cdots, k$, $\rho_i(x,t)$ satisfies (1.11) if and only if

$$u_i(x,t) = \exp\left[h^T(x,t) S^{-1}(t) y_{\tau_{i-1}}\right] \rho_i(x,t), \tag{1.12}$$

satisfies the Kolmogorov forward equation

$$\frac{\partial u_i}{\partial t}(x,t) = \left(L - \frac{1}{2} h^T S^{-1} h\right) u_i(x,t), \tag{1.13}$$

where L is defined in (1.3).

To be more specific, how to shift the computations off-line. Let us consider the initial data to Kolmogorov forward equation (1.13) as a set of complete orthonormal base in $L^2(\mathbb{R}^n)$, say $\{\phi_i(x)\}_{i=1}^{\infty}$. For arbitrary initial condition $u_i(x,\tau_{i-1}) \in L^2(\mathbb{R}^n)$, it can be represented as $u_i(x,\tau_{i-1}) = \sum_{n=1}^{\infty} \hat{u}_{i,n}\phi_n(x)$. Suppose the operator $(L - \frac{1}{2}h^T S^{-1}h)$ generates a C_0-semigroup (under some conditions), denoted as $\mathcal{S}(t)$. Hence, the solution to (1.13) with the initial condition $u_i(x,\tau_{i-1})$ can be expressed as

$$u_i(x,\tau_i) = \mathcal{S}(\tau_i - \tau_{i-1})u_i(x,\tau_{i-1}) = \sum_{n=1}^{\infty} \hat{u}_{i,n}\mathcal{S}(\tau_i - \tau_{i-1})\phi_n(x),$$

where $\mathcal{S}(\tau_i - \tau_{i-1})\phi_n(x)$ can be computed off-line.

Next, we specify the observation updates (the initial condition of (1.13)) for each time step. For $0 \le t \le \tau_1$, the initial condition is $u_1(x,0) = \sigma_0(x)$. At time $t = \tau_1$,

$$\rho_1(x,\tau_1) = \exp\left[-h^T(x,\tau_1)S^{-1}(\tau_1)y_0\right]u_1(x,\tau_1),$$

by (1.12). Combined with (1.11), we have

$$\rho_2(x,\tau_1) = \exp\left[-h^T(x,\tau_1)S^{-1}(\tau_1)y_0\right]u_1(x,\tau_1).$$

Now that the observation y_{τ_1} is available, the prediction $u_1(x,\tau_1)$ can be updated as:

$$u_2(x,\tau_1) = \exp[h^T(x,\tau_1)S^{-1}(\tau_1)y_{\tau_1}]\rho_2(x,\tau_1) = \exp[h^T(x,\tau_1)S^{-1}(\tau_1)y_{\tau_1}]u_1(x,\tau_1).$$

This is the initial condition in the next time interval $\tau_1 \le t \le \tau_2$. Recursively, the initial condition for $\tau_{i-1} \le t \le \tau_i$ is

$$u_i(x,\tau_{i-1}) = \exp[h^T(x,\tau_{i-1})S^{-1}(\tau_{i-1})(y_{\tau_{i-1}} - y_{\tau_{i-2}})]u_{i-1}(x,\tau_{i-1}), \tag{1.14}$$

for $i = 2, 3, \cdots, k$. The approximation of $\rho(x,t)$, denoted as $\hat{\rho}(x,t)$, is obtained

$$\hat{\rho}(x,t) = \sum_{i=1}^{k} \chi_{[\tau_{i-1},\tau_i]}(t)\rho_i(x,t), \tag{1.15}$$

where $\rho_i(x,t)$ is obtained by (1.12). Furthermore, we could recover the approximation of $\sigma(x,t)$ by (1.4). Till now, we have briefly described Yau-Yau's algorithm. A natural question comes to us: If we applied this algorithm to the most general setting of DMZ equation (1.5) or (1.6), then

> *is* $\hat{\rho}(x,t)$ *a good approximation of* $\rho(x,t)$*, for* $(x,t) \in \mathbb{R}^n \times [0,T]$*, as* $|\mathcal{P}_k| \to 0$*? If it is, then in what sense?*

To answer this question, we divide it into two steps: first, we show that the density restricted in a bounded domain $\Omega \in \mathbb{R}^n$, denoted as ρ_Ω, can be well approximated by gluing up the piecewise approximation $\hat{\rho}_\Omega$, similarly obtained as (1.15); next, it is left to show that ρ_Ω is a good approximation of ρ in $\mathbb{R}^n$.

The first step is accomplished by the theorem below:

Theorem 1.1: Let Ω be a bounded domain in $\mathbb{R}^n$. Assume that

1) $|N(x,t)| \leq C,$ (1.16)

2) There exists some $\alpha \in (0,1)$, such that

$$|N(x,t) - N(x,t;\bar{t})| \leq \tilde{C}|t-\bar{t}|^\alpha, \tag{1.17}$$

for all $(x,t) \in \Omega \times [0,T]$, $\bar{t} \in [0,T]$, where

$$N(x,t) \equiv -\frac{\partial}{\partial t}\left(h^T S^{-1}\right) y_t - \frac{1}{2}D_w^2 K + \frac{1}{2}D_w K \cdot \nabla K - f \cdot \nabla K - \frac{1}{2}\left(h^T S^{-1} h\right), \tag{1.18}$$

and $N(x,t;\bar{t})$ denotes the observation y_t contained in $N(x,t)$ be freezed at $y_{\bar{t}}$. Let $\rho_\Omega(x,t)$ be the solution of (1.6) on $\Omega \times [0,T]$:

$$\begin{cases} \dfrac{\partial \rho_\Omega}{\partial t}(x,t) = \dfrac{1}{2}D_w^2 \rho_\Omega(x,t) + F(x,t)\cdot\nabla\rho_\Omega(x,t) + J(x,t)\rho_\Omega(x,t) \\ \rho_\Omega(x,0) = \sigma_{0,\Omega}(x) \\ \rho_\Omega(x,t)|_{\partial\Omega} = 0, \end{cases} \tag{1.19}$$

where D_w^2, $F(x,t)$ and $J(x,t)$ are defined in (1.7)-(1.9) and $\sigma_{0,\Omega}$ is defined as

$$\sigma_{0,\Omega}(x) = \begin{cases} \sigma_0(x), & x \in \Omega_\epsilon \\ 0, & x \in \mathbb{R}^n \setminus \Omega, \end{cases} \tag{1.20}$$

where $\Omega_\epsilon = \{x \in \Omega : \ \mathrm{dist}(x, \partial\Omega) > \epsilon\}$.

For any $0 \leq \tau \leq T$, let $\mathcal{P}_k = \{0 = \tau_0 < \tau_1 < \tau_2 < \cdots < \tau_k = \tau\}$ be a partition of $[0,\tau]$, where $\tau_i = \frac{i\tau}{k}$. Let $\rho_{i,\Omega}(x,t)$ be the solution on $\Omega \times [\tau_{i-1}, \tau_i]$ of the equation

$$\begin{cases} \dfrac{\partial \rho_{i,\Omega}}{\partial t}(x,t) = \dfrac{1}{2}D_w^2 \rho_{i,\Omega}(x,t) + F(x,t;\tau_{i-1})\cdot\nabla\rho_{i,\Omega}(x,t) + J(x,t;\tau_{i-1})\rho_{i,\Omega}(x,t) \\ \rho_{i,\Omega}(x,\tau_{i-1}) = \rho_{i-1,\Omega}(x,\tau_{i-1}) \\ \rho_{i,\Omega}(x,t)|_{\partial\Omega} = 0, \end{cases} \tag{1.21}$$

for $i = 1, 2, \cdots, k$, with the convention that $\rho_{1,\Omega}(x,0) = \sigma_{0,\Omega}(x)$. Then

$$\rho_\Omega(x,\tau) = \lim_{k\to\infty} \rho_{k,\Omega}(x,\tau),$$

in the L^1 sense in space and the following estimate holds:

$$\int_\Omega |\rho_\Omega - \rho_{k,\Omega}|(x,\tau) \leq C\frac{1}{k^\alpha}, \tag{1.22}$$

where $C = C(T, \int_\Omega \sigma_{0,\Omega})$ is a generic constant. The right-hand side of (1.22) tends to zero as $k \to \infty$.

The following theorem verifies that ρ_Ω, $\Omega = B_R$, is a good approximation of ρ, provided that R is sufficiently large.

Theorem 1.2: For any $T > 0$, let $\rho(x,t)$ be a solution of the robust DMZ equation (1.6) in $\mathbb{R}^n \times [0,T]$. Assume the following conditions are satisfied, for all $(x,t) \in \mathbb{R}^n \times [0,T]$:

1) $N(x,t) + \frac{3}{2}n\left|\left|GQG^T\right|\right|_\infty + |f - D_wK| \leq C,$ (1.23)

2) $e^{-\sqrt{1+|x|^2}}\left[14n\left|\left|GQG^T\right|\right|_\infty + 4\,|f - D_wK|\right] \leq \tilde{C},$ (1.24)

where C, $\tilde{C}$ are constants possibly depending on T. Let $R \gg 1$ and ρ_R be the solution to the robust DMZ equation (1.6) on the ball B_R:

$$\begin{cases} \dfrac{\partial \rho_R}{\partial t}(x,t) = \dfrac{1}{2}D_w^2\rho_R(x,t) + F(x,t)\nabla\rho_R(x,t) + J(x,t)\rho_R(x,t) \\ \rho_R(x,0) = \sigma_{0,R}(x) \\ \rho_R(x,t) = 0 \qquad \text{for } (x,t) \in \partial B_R \times [0,T], \end{cases} \tag{1.25}$$

where $F(x,t)$, $J(x,t)$ and $\sigma_{0,R}$ are defined as (1.8), (1.9) and (1.20), respectively. Let $v = \rho - \rho_R$. Then $v \geq 0$ for all $(x,t) \in B_R \times [0,T]$ and

$$\int_{B_R} \phi v(x,T) \leq Ce^{-R}\int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}}\sigma_0(x), \tag{1.26}$$

where $C = C(T)$ is a generic constant and $\phi(x) = e^{-R[(|x|^2/R^2-1)^2+1]} - e^{-R}$. In particular,

$$\int_{B_{\frac{R}{2}}} v(x,T) \leq Ce^{-\frac{9}{16}R}\int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}}\sigma_0(x), \tag{1.27}$$

where $C = C(T)$.

Also, some interesting properties of the density function have been discovered. For example, the growth rate of the density function in a fixed large ball with respect to time (Proposition 3.2); the decaying rate of the density outside a large ball with respect to the radius (Theorem 3.4); the lower bound of the density function inside a large ball in terms of time (Theorem 5.5), etc.

This paper is organized as following: section II is the preliminary and the statement of the well-posedness of the robust DMZ equation, whose proof is left in appendix; the proof of Theorem 1.2 and Theorem 3.4 are presented in section III; section IV is devoted to the proof of Theorem 1.1; Theorem 5.5 can be found in section V and the numerical simulations of 1D cubic sensor are illustrated in section VI.

II. PRELIMINARY

Let $\mathbb{Q}_T = \mathbb{R}^n \times [0, T]$. Assume that the following conditions hold for the entire paper:

(1) The operator L defined in (1.3) is a uniform elliptic operator and it is bounded from above on $\mathbb{Q}_T$, i.e. there exists constants $\lambda > 0$ such that

$$\lambda|\xi|^2 \leq \sum_{i,j=1}^{n}(GQG^T)_{ij}\xi_i\xi_j,$$

for any $(x,t) \in \mathbb{Q}_T$, for any $\xi = (\xi_1, \xi_2, \cdots, \xi_n) \in \mathbb{R}^n$. And

$$||GQG^T||_\infty = \sup_{(x,t)\in\mathbb{Q}_T}|GQG^T|_\infty < \infty,$$

where $|\cdot|_\infty$ is the sup-norm of the matrix.

(2) The initial density function $\sigma_0(x) \in H^1(\mathbb{R}^n)$ decays fast enough. To be more specific, we require that

$$\int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}}\sigma_0(x) < \infty.$$

A. *Existence and uniqueness of the nonnegative weak solution*

Let $H^1(\mathbb{R}^n)$ be the space of the functions with their gradient and themselves are square integrable. It equips the norm

$$||u(x)||_1^2 = \int_{\mathbb{R}^n}(u^2 + |\nabla_x u|^2)dx.$$

Let $H^{1;1}(\mathbb{Q}_T)$ be the subspace of $H^1(\mathbb{R}^n)$ for each time $t \geq 0$ and its derivative with respect to t is also square integrable. The norm is defined as

$$||v(x,t)||_{1;1}^2 = \int_{\mathbb{Q}_T}(v^2 + |\nabla_x v|^2 + |\partial_t v|^2)dxdt.$$

It is well known [8] that $H^1(\mathbb{R}^n)$ and $H^{1;1}(\mathbb{Q}_T)$ are complete. It can be shown that under some mild conditions, the existence and uniqueness of a nonnegative weak solution to the DMZ equation (1.6) on $\mathbb{R}^n \times [0,T]$ have been guaranteed. We state the theorems here and leave the proofs in the Appendix.

Theorem 2.3: Under the condition (1)-(2) and the condition (A.1)-(A.5) in Theorem A.1, the robust DMZ equation (1.6) on $\mathbb{Q}_T$ with the initial value $\sigma_0 \in H^1(\mathbb{R}^n)$ admits a nonnegative weak solution. Assume further that for some $c > 0$,

$$\sup_{0\le t\le T}\int_{\mathbb{R}^n} e^{cr}\rho^2(x,t)dx < \infty,$$

and

$$\int_{\mathbb{Q}_T} |\nabla\rho(x,t)|^2 dxdt < \infty,$$

where $r = |x|$. Suppose that there exists a finite number $\alpha > 0$ such that

$$2J(x,t) - \frac{1}{4\lambda_1}[cD_w r - (F(x,t) + \tilde{F}(x,t))]^2 \le \alpha, \tag{2.1}$$

for all $(x,t) \in \mathbb{Q}_T$, where λ_1 is the smallest eigenvalue of the matrix (GQG^T),

$$\tilde{F}(x,t) = \left[\frac{1}{2}\sum_{j=1}^{n}(GQG^T)_{ij} + \sum_{j=1}^{n}(GQG^T)_{ij}\frac{\partial K}{\partial x_j} - f_i\right]_{i=1}^{n}, \tag{2.2}$$

and $J(x,t)$ is defined as in (1.9). Then the weak solution $\rho(x,t)$ of the robust DMZ equation on $\mathbb{Q}_T$ is unique.

B. Technical lemma

In the proofs of the theorems, we will repeatedly adopt the following lemma with suitable chosen test functions. We'd like to prove the lemma here.

Lemma 2.1: Assume that ρ_Ω satisfies the robust DMZ equation (1.6) on some bounded domain $\Omega \in \mathbb{R}^n$, for $0 \le t \le T$. Then, for any test function $\psi(x) \in C^\infty(\Omega)$ (for concise notation, we omit the subscript Ω below), we have

$$\begin{aligned}\frac{d}{dt}\int_\Omega \psi\rho =& \frac{1}{2}\int_\Omega D_w^2\psi\rho + \int_\Omega (f - D_w K)\cdot\nabla\psi\rho + \int_\Omega \psi\rho N \\ &+ \frac{1}{2}\int_{\partial\Omega}\psi\left(D_w\rho\cdot\nu\right) - \frac{1}{2}\int_{\partial\Omega}\rho\left(D_w\psi\cdot\nu\right) + \frac{1}{2}\int_{\partial\Omega}\psi\rho\sum_{i,j=1}^{n}\frac{\partial}{\partial x_i}\left(GQG^T\right)_{ij}\nu_j \\ &+ \int_{\partial\Omega}\psi\rho\left(D_w K\cdot\nu\right) - \int_{\partial\Omega}\psi\rho(f\cdot\nu),\end{aligned} \tag{2.3}$$

where $\nu = (\nu_1, \nu_2, \cdots, \nu_n)$ is the exterior normal vector of Ω,

$$D_w * = \left[\sum_{j=1}^{n} \left(GQG^T\right)_{ij} (x,t) \frac{\partial *}{\partial x_j} \right]_{i=1}^{n}, \tag{2.4}$$

and D_w^2, N and K are defined in (1.7), (1.18) and (1.10), respectively.

Remark 2.1: We don't specify the test function $\psi(x)$ on the boundary $\partial\Omega$. All the boundary terms in (2.3) are kept. However, in the proofs of the theorems, all the boundary terms are vanishing by choosing the test function ψ in Lemma 2.1 properly.

Proof: Multiplying $\psi(x)$ on both sides of the robust DMZ equation (1.6) and integrating over the domain Ω yields

$$\frac{d}{dt} \int_\Omega \psi\rho = \int_\Omega \psi \left[\frac{1}{2} D_w^2 \rho + F(x,t) \cdot \nabla\rho + J(x,t)\rho \right], \tag{2.5}$$

where $F(x,t)$ and $J(x,t)$ are defined in (1.8) and (1.9), respectively. Apply integration by parts to the first two terms on the right-hand side of (2.5):

$$\begin{aligned}
\int_\Omega \psi D_w^2 \rho =& \int_\Omega \sum_{i,j=1}^{n} \frac{\partial^2 \psi}{\partial x_i \partial x_j} \left(GQG^T\right)_{ij} \rho + 2 \int_\Omega \sum_{i,j=1}^{n} \frac{\partial \psi}{\partial x_i} \frac{\partial}{\partial x_j} \left(GQG^T\right)_{ij} \rho \\
&+ \int_\Omega \sum_{i,j=1}^{n} \frac{\partial^2}{\partial x_i \partial x_j} \left(GQG^T\right)_{ij} \psi\rho \\
&+ \int_{\partial\Omega} \psi \sum_{i,j=1}^{n} \left(GQG^T\right)_{ij} \frac{\partial \rho}{\partial x_j} \nu_i - \int_{\partial\Omega} \rho \sum_{i,j=1}^{n} \frac{\partial \psi}{\partial x_i} \left(GQG^T\right)_{ij} \nu_j \\
&- \int_{\partial\Omega} \rho\psi \sum_{i,j=1}^{n} \frac{\partial}{\partial x_i} \left(GQG^T\right)_{ij} \nu_j,
\end{aligned} \tag{2.6}$$

$$\begin{aligned}
\int_\Omega \psi F(x,t) \cdot \nabla\rho =& - \int_\Omega \sum_{i,j=1}^{n} \frac{\partial \psi}{\partial x_i} \frac{\partial}{\partial x_j} \left(GQG^T\right)_{ij} \rho - \int_\Omega \sum_{i,j=1}^{n} \frac{\partial^2}{\partial x_i \partial x_j} \left(GQG^T\right)_{ij} \psi\rho \\
&- \int_\Omega \sum_{i,j=1}^{n} \frac{\partial \psi}{\partial x_i} \left(GQG^T\right)_{ij} \frac{\partial K}{\partial x_j} \rho - \int_\Omega \psi\rho \sum_{i,j=1}^{n} \frac{\partial}{\partial x_i} \left(GQG^T\right)_{ij} \frac{\partial K}{\partial x_j} \\
&- \int_\Omega \psi\rho \sum_{i,j=1}^{n} \left(GQG^T\right)_{ij} \frac{\partial^2 K}{\partial x_i \partial x_j} + \int_\Omega \psi\rho \sum_{i=1}^{n} \frac{\partial f_i}{\partial x_i} + \int_\Omega \rho \sum_{i=1}^{n} \frac{\partial \psi}{\partial x_i} f_i \\
&+ \int_{\partial\Omega} \psi\rho \sum_{i,j=1}^{n} \frac{\partial}{\partial x_j} \left(GQG^T\right)_{ij} \nu_i + \int_{\partial\Omega} \psi\rho \sum_{i,j=1}^{n} \left(GQG^T\right)_{ij} \frac{\partial K}{\partial x_j} \nu_i \\
&- \int_{\partial\Omega} \psi\rho \sum_{i=1}^{n} f_i \nu_i.
\end{aligned} \tag{2.7}$$

Substituting (2.6) and (2.7) back into (2.5), (2.3) is obtained by written in compact notations.□

III. REDUCTION OF THE PROBLEM TO THE BOUNDED DOMAIN CASE

In this section we shall prove that the solution ρ to the robust DMZ equation (1.6) in $\mathbb{R}^n$ can be well approximated by the solution ρ_R of (1.25) in a large ball B_R. Moreover, the error estimate with respect to the radius R is given explicitly in the L^1 sense. Let C, $\tilde{C}$ and $\hat{C}$ denote the generic constants, which may differ from line to line.

We first show an interesting proposition, which reflects how the density function in the large ball changing with respect to time. It is also an important ingredient in estimating the error.

Proposition 3.2: For any $T > 0$, let $\rho_R(x,t)$ be a solution of the robust DMZ equation (1.19) with $\Omega = B_R$ and $0 \le t \le T$. Assume that condition (1.23) is satisfied. Then

$$\int_{B_R} e^{\sqrt{1+|x|^2}} \rho_R(x,t) \le e^{Ct} \int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}} \sigma_0(x). \tag{3.1}$$

Proof: Choose the test function in Lemma 2.1 $\psi = e^{\phi_1}$, where $\phi_1 \in C^\infty(B_R)$, $B_R = \{x \in \mathbb{R}^n : |x| \le R\}$. Let ρ_R be the solution of the robust DMZ equation (1.25) on the ball B_R. By Lemma 2.1, we have

$$\begin{aligned} \frac{d}{dt} \int_{B_R} e^{\phi_1} \rho_R = &\int_{B_R} e^{\phi_1} \rho_R \left[\frac{1}{2} \left(D_w^2 \phi_1 + D_w \phi_1 \cdot \nabla \phi_1 \right) + (f - D_w K) \cdot \nabla \phi_1 + N \right] \\ &+ \frac{1}{2} \int_{\partial B_R} e^{\phi_1} (D_w \rho_R \cdot \nu). \end{aligned} \tag{3.2}$$

All the boundary integrations in (2.3), except the first term, vanishes, since $\rho_R|_{\partial\Omega} = 0$. Moreover, recall that $\rho_R \ge 0$ in B_R and vanishes on ∂B_R implies that $\frac{\partial \rho_R}{\partial \nu}|_{\partial B_R} \le 0$. Hence, on ∂B_R,

$$(D_w \rho_R \cdot \nu) = \sum_{i=1}^n \left[\sum_{j=1}^n (GQG^T)_{ij} \frac{\partial \rho_R}{\partial r} \frac{\partial r}{\partial x_j} \right] \nu_i = \frac{\partial \rho_R}{\partial r} \left[\sum_{i,j=1}^n (GQG^T)_{ij} \frac{x_j}{r} \frac{x_i}{r} \right] \le 0,$$

by the positive definite assumption of (GQG^T). Thus, (3.2) can be reduced further

$$\frac{d}{dt} \int_{B_R} e^{\phi_1} \rho_R \le \int_{B_R} e^{\phi_1} \rho_R \left[\frac{1}{2} \left(D_w^2 \phi_1 + D_w \phi_1 \cdot \nabla \phi_1 \right) + (f - D_w K) \cdot \nabla \phi_1 + N \right]. \tag{3.3}$$

Choose $\phi_1(x) = \sqrt{1+|x|^2}$ and estimate the terms containing ϕ_1 on the right-hand side of (3.2) one by one:

$$\begin{aligned} D_w^2 \phi_1 &= \sum_{i=1}^n \left(GQG^T \right)_{ii} \frac{1}{\sqrt{1+|x|^2}} - \sum_{i,j=1}^n \left(GQG^T \right)_{ij} \frac{x_i x_j}{(1+|x|^2)^{\frac{3}{2}}} \\ &\le \left|\left| GQG^T \right|\right|_\infty \left[\frac{n}{\sqrt{1+|x|^2}} + \frac{n|x|^2}{(1+|x|^2)^{\frac{3}{2}}} \right] \le 2n \left|\left| GQG^T \right|\right|_\infty, \end{aligned} \tag{3.4}$$

$$D_w \phi_1 \cdot \nabla \phi_1 = \sum_{i,j=1}^n \left(GQG^T \right)_{ij} \frac{x_i x_j}{1+|x|^2} \le \left|\left| GQG^T \right|\right|_\infty \frac{\sum_{i,j=1}^n x_i x_j}{1+|x|^2} \le n \left|\left| GQG^T \right|\right|_\infty, \tag{3.5}$$

and

$$|(f - D_w K) \cdot \nabla \phi_1| \le |f - D_w K| \cdot \frac{|x|}{\sqrt{1+|x|^2}} \le |f - D_w K|, \tag{3.6}$$

where $|\cdot|$ is the Euclidean norm. Substitute the estimate (3.4)-(3.6) back into (3.3), we get

$$\frac{d}{dt}\int_{B_R} e^{\phi_1}\rho_R \le \int_{B_R} e^{\phi_1}\rho_R \left[\frac{3}{2}n \left|\left|GQG^T\right|\right|_\infty + |f - D_w K| + N\right] \le C\int_{B_R} e^{\phi_1}\rho_R,$$

by condition (1.23). Hence,

$$\int_{B_R} e^{\phi_1}\rho_R(x,t) \le e^{Ct}\int_{B_R} e^{\phi_1}\rho_R(x,0) \le e^{Ct}\int_{\mathbb{R}^n} e^{\phi_1}\rho(x,0) = e^{Ct}\int_{\mathbb{R}^n} e^{\phi_1}\sigma_0(x),$$

for $0 \le t \le T$. □

We are ready to show that the solution ρ_R to (1.25) on B_R is a good approximation of the solution ρ on $\mathbb{R}^n$.

Proof of Theorem 1.2: By the maximum principle (cf. Theorem 1, p. 34 in Friedman's book [4]), we have $v = \rho - \rho_R \ge 0$ for $(x,t) \in B_R \times [0,T]$, since $v|_{\partial B_R} \ge 0$ for $0 \le t \le T$. Let ψ in Lemma 2.1 as

$$\varrho(x) = e^{-\phi_2(x)} - e^{-R},$$

where ϕ_2 is a radial symmetric function such that $\phi_2(x)|_{\partial B_R} = R$, $\nabla\phi_2|_{\partial B_R} = 0$ and ϕ_2 is increasing in $|x|$. Hence, $\varrho|_{\partial B_R} = 0$ and $\nabla\varrho|_{\partial B_R} = 0$. Apply Lemma 2.1 to v, taking the place of ρ_R, with the test function $\psi = \varrho$, we have

$$\begin{aligned}
\frac{d}{dt}\int_{B_R} \varrho v &= \int_{B_R} v\left[\frac{1}{2}D_w^2\varrho + (f - D_w K)\cdot\nabla\varrho + \varrho N\right] \\
&= \int_{B_R} v\left\{\frac{1}{2}e^{-\phi_2}\left(D_w\phi_2\cdot\nabla\phi_2 - D_w^2\phi_2\right) - e^{-\phi_2}\left(f - D_w K\right)\cdot\nabla\phi_2 + \varrho N\right\} \\
&= \int_{B_R} v\varrho\left[-\frac{1}{2}D_w^2\phi_2 + \frac{1}{2}D_w\phi_2\cdot\nabla\phi_2 - (f - D_w K)\cdot\nabla\phi_2 + N\right] \\
&\quad + e^{-R}\int_{B_R} e^{\sqrt{1+|x|^2}} v\left[e^{-\sqrt{1+|x|^2}}\left(-\frac{1}{2}D_w^2\phi_2 + \frac{1}{2}D_w\phi_2\cdot\nabla\phi_2 - (f - D_w K)\cdot\nabla\phi_2\right)\right] \\
&\triangleq \int_{B_R} v\varrho \mathrm{I}_1 + e^{-R}\int_{B_R} e^{\sqrt{1+|x|^2}} v \mathrm{I}_2.
\end{aligned}$$

Let us choose $\phi_2(x)$ to be $\phi_2(x) = R\vartheta(\frac{|x|^2}{R^2})$, where $\vartheta(x) = 1 - (1-x)^2$. It is easy to check that $\phi_2(x)$ satisfies all the conditions we mentioned before. Direct computations yield, for any

$x \in B_R$, $R >> 1$,

$$\begin{aligned}\left|D_w^2\phi_2\right| &= \left|\sum_{i,j=1}^n \left(GQG^T\right)_{ij}\left(-\frac{8x_ix_j}{R^3}\right) + \sum_{i=1}^n \left(GQG^T\right)_{ii}\frac{4}{R}\left(1-\frac{|x|^2}{R^2}\right)\right| \\ &\le \left|\left|GQG^T\right|\right|_\infty \left(\frac{8n|x|^2}{R^3}+\frac{4n}{R}\right) \le 12n\left|\left|GQG^T\right|\right|_\infty,\end{aligned} \tag{3.7}$$

$$|D_w\phi_2\cdot\nabla\phi_2| = \left|\left(1-\frac{|x|^2}{R^2}\right)^2\sum_{i,j=1}^n\left(GQG^T\right)_{ij}\frac{4x_i}{R}\frac{4x_j}{R}\right| \le 16n\left|\left|GQG^T\right|\right|_\infty, \tag{3.8}$$

and

$$|(f-D_wK)\cdot\nabla\phi_2| = \left|(f-D_wK)\frac{4x}{R}\left(1-\frac{|x|^2}{R^2}\right)\right| \le 4\left|f-D_wK\right|. \tag{3.9}$$

It follows that

$$\sup_{B_R}|\mathrm{I}_1| \le 14n\left|\left|GQG^T\right|\right|_\infty + 4\left|f-D_wK\right| + N \le C,$$

by condition (1.23). Similarly,

$$\sup_{B_R}|\mathrm{I}_2| \le \sup_{B_R}\left[e^{-\sqrt{1+|x|^2}}\left(14n\left|\left|GQG^T\right|\right|_\infty + 4\left|f-D_wK\right|\right)\right] \le \tilde{C},$$

by condition (1.24). In the view of Proposition 3.2, one gets

$$\frac{d}{dt}\int_{B_R}\varrho v \le C\int_{B_R}\varrho v + e^{-R}\tilde{C}\int_{B_R}e^{\sqrt{1+|x|^2}}\rho \le C\int_{B_R}\varrho v + e^{-R+\hat{C}T}\tilde{C}\int_{\mathbb{R}^n}e^{\sqrt{1+|x|^2}}\sigma_0(x). \tag{3.10}$$

Multiply e^{-Ct} on both sides of (3.10) yields

$$\frac{d}{dt}\left[e^{-Ct}\int_{B_R}\varrho v\right] \le e^{-R+\hat{C}T-Ct}\tilde{C}\int_{\mathbb{R}^n}e^{\sqrt{1+|x|^2}}\sigma_0(x).$$

Integrate from 0 to T and multiply e^{CT} on both sides gives us

$$\int_{B_R}\varrho v(x,T) \le ||v(x,0)||_\infty e^{CT}\int_{B_R}\varrho dx + \frac{e^{CT}-1}{C}e^{-R+\hat{C}T}\tilde{C}\int_{\mathbb{R}^n}e^{\sqrt{1+|x|^2}}\sigma_0(x),$$

where $v(x,0)=\sigma_0-\sigma_{0,R}$. Recall that $\varrho(x) = e^{-R\left[-\left(|x|^2/R^2-1\right)^2+1\right]} - e^{-R}$, $|x|\le R$, we estimate

$$\int_{B_R}\varrho \le \int_{B_R}\left(1-e^{-R}\right) \le CR^n$$

and

$$\int_{B_R}\varrho v(x,T) \ge \int_{B_{\frac{R}{2}}}\left(e^{-R\left[-\left(|x|^2/R^2-1\right)^2+1\right]} - e^{-R}\right)v(x,T) \ge \frac{1}{2}e^{-\frac{7}{16}R}\int_{B_{\frac{R}{2}}}v(x,T).$$

It is easy to see that $||v(x,0)||_\infty \int_{B_R} \varrho \leq C(n)\epsilon R^n$ is arbitrarily small, since ϵ is independent of R. It yields that

$$\int_{B_{\frac{R}{2}}} v(x,T) \leq Ce^{-\frac{9}{16}R} \int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}}\sigma_0(x), \tag{1.28}$$

where C is a generic constant, which depends on T. □

By refining the proofs of Proposition 3.2 and Theorem 1.2, we obtain an interesting property of the density function $\rho(x,t)$. It asserts that ρ captures almost all the density in a large ball. And we could give an precise estimate of the density outside the large ball.

Theorem 3.4: Let $\rho(x,t)$ be a solution of the robust DMZ equation (1.6) in $\mathbb{Q}_T$. Assume that

1) Condition (1.23) is satisfied;
2) A stronger version of condition (1.24) is valid. To be more precise,

$$e^{-\frac{1}{2}\sqrt{1+|x|^2}} \left[16n \left|\left|GQG^T\right|\right|_\infty + 4\left|f - D_w K\right|\right] \leq C, \tag{3.11}$$

   for all $(x,t) \in \mathbb{Q}_T$.

Then

$$\int_{|x|\geq R} \rho(x,T) \leq Ce^{-\frac{1}{2}\sqrt{1+R^2}} \int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}}\sigma_0(x), \tag{3.12}$$

where C is a generic constant, which depends on T.

Proof: Let $v = \rho - \rho_R$ as in the proof of Theorem 1.2. By the maximum principle, we have that $v \geq 0$ for all $(x,t) \in B_R \times [0,T]$. Choose the test function ψ in Lemma 2.1 as

$$\Phi(x) = \gamma(x)\varrho(x),$$

where $\gamma(x) = e^{\frac{1}{2}\phi_1(x)}$ and $\phi_1(x)$, $\varrho(x)$ are defined in the proof of Proposition 3.2 and Theorem 1.2. It follows directly that $\Phi|_{\partial B_R} = \nabla_x \Phi|_{\partial B_R} = 0$, by the fact that $\varrho|_{\partial B_R} = \nabla\varrho|_{\partial B_R} = 0$. Apply Lemma 2.1 to v taking place of ρ_R with the test function Φ, we have

$$\begin{aligned}\frac{d}{dt}\int_{B_R} \Phi v &= \frac{1}{2}\int_{B_R} D_w^2 \Phi v + \int_{B_R} (f - D_w K)\cdot \Phi v + \int_{B_R} \Phi N v \\ &= \frac{1}{2}\int_{B_R} (D_w^2\gamma\varrho + 2D_w\gamma \cdot \nabla\varrho + \gamma D_w^2\varrho)v + \int_{B_R}(f - D_w K)\cdot(\nabla\gamma\varrho + \gamma\nabla\varrho)v \\ &\quad + \int_{B_R} \gamma\varrho N v.\end{aligned}$$

All the boundary integrations vanish due to the similar arguments in Theorem 1.2. Direct computations with $\gamma(x) = e^{\frac{1}{2}\phi_1(x)}$ and $\varrho(x) = e^{-\phi_2(x)} - e^{-R}$ by hand give us

$$\begin{aligned}
\frac{d}{dt}\int_{B_R} \Phi v =& \frac{1}{2}\int_{B_R} \left[\frac{1}{2}e^{\frac{1}{2}\phi_1}\left(D_w^2\phi_1 + \frac{1}{2}D_w\phi_1 \cdot \nabla\phi_1\right)\varrho - e^{\frac{1}{2}\phi_1}D_w\phi_1 \cdot e^{-\phi_2}\nabla\phi_2\right. \\
&\qquad\qquad \left. + \gamma e^{-\phi_2}\left(D_w\phi_2 \cdot \nabla\phi_2 - D_w^2\phi_2\right)\right] v \\
&+ \int_{B_R} (f - D_wK) \cdot \left(\frac{1}{2}e^{\frac{1}{2}\phi_1}\nabla\phi_1\varrho - \gamma e^{-\phi_2}\nabla\phi_2\right) v + \int_{B_R} \gamma\varrho N v \\
=& \int_{B_R} \Phi v \left[\frac{1}{4}\left(D_w^2\phi_1 + \frac{1}{2}D_w\phi_1 \cdot \nabla\phi_1\right) - \frac{1}{2}D_w\phi_1 \cdot \nabla\phi_2 + \frac{1}{2}\left(D_w\phi_2 \cdot \nabla\phi_2 - D_w^2\phi_2\right)\right. \\
&\qquad\qquad \left. + (f - D_wK) \cdot (\frac{1}{2}\nabla\phi_1 - \nabla\phi_2) + N\right] \\
&+ e^{-R}\int_{B_R} \gamma v \left[-\frac{1}{2}D_w\phi_1 \cdot \nabla\phi_2 + \frac{1}{2}\left(D_w\phi_2 \cdot \nabla\phi_2 - D_w^2\phi_2\right) - (f - D_wK) \cdot \nabla\phi_2\right] \\
\triangleq& \int_{B_R} \Phi v[\mathrm{I}_3] + e^{-R}\int_{B_R} \gamma v[\mathrm{I}_4],
\end{aligned}$$

By the similar estimates (3.4)-(3.6), (3.7)-(3.9), we have

$$\begin{aligned}
\sup_{B_R} |\mathrm{I}_3| &\le 17n \left|\left|GQG^T\right|\right|_\infty + 5|f - D_wK| + N, \\
\sup_{B_R} |\mathrm{I}_4| &\le 16n \left|\left|GQG^T\right|\right|_\infty + 4|f - D_wK|.
\end{aligned}$$

Hence,

$$\begin{aligned}
\frac{d}{dt}\int_{B_R} \Phi v \le& C\int_{B_R} \Phi v + e^{-R}\tilde{C}\int_{B_R} e^{\phi_1} v \le C\int_{B_R} \Phi v + e^{-R}\tilde{C}\int_{B_R} e^{\phi_1}\rho \\
\le& C\int_{B_R} \Phi v + \tilde{C}e^{-R+Ct}\int_{B_R} e^{\phi_1}\sigma_0(x) \le C\int_{B_R} \Phi v + \tilde{C}e^{-R+Ct}\int_{\mathbb{R}^n} e^{\phi_1}\sigma_0(x),
\end{aligned}$$

by condition (1.23), (1.24) and (3.1). By the similar argument in the proof of Theorem 1.2, where we get the estimate of $\int_{B_R} \varrho v$, we have

$$\int_{B_R} \Phi v(x, T) \le Ce^{-R}\int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}}\sigma_0(x), \tag{3.13}$$

where C is a generic constant, which depends on T. Recall that $\varrho(x) = e^{-R[-(|x|^2/R^2-1)^2+1]} - e^{-R}$, it implies that

$$\int_{B_R} \Phi v(x, T) \ge \frac{1}{2}e^{-\frac{7}{16}R}\int_{B_{\frac{R}{2}}} \gamma v(x, T). \tag{3.14}$$

Combine (3.13) and (3.14), we obtain that

$$\int_{B_{\frac{R}{2}}} \gamma v(x,T) \le Ce^{-\frac{9}{16}R} \int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}} \sigma_0(x).$$

This implies that

$$\int_{B_{\frac{R}{2}}} \gamma \rho(x,T) \le \int_{B_{\frac{R}{2}}} \gamma \rho_R(x,T) + Ce^{-\frac{9}{16}R} \int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}} \sigma_0(x) \le C(1+e^{-\frac{9}{16}R}) \int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}} \sigma_0(x),$$

by (3.1). Let $R \to \infty$,

$$\int_{\mathbb{R}^n} \gamma \rho(x,T) \le C \int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}} \sigma_0(x).$$

Consider the integration outside the large ball B_R,

$$e^{\frac{1}{2}\sqrt{1+R^2}} \int_{|x|\ge R} \rho(x,T) \le \int_{|x|\ge R} \gamma \rho(x,T) \le C \int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}} \sigma_0(x).$$

Therefore, we achieve the conclusion that

$$\int_{|x|\ge R} \rho(x,T) \le Ce^{-\frac{1}{2}\sqrt{1+|R|^2}} \int_{\mathbb{R}^n} e^{\sqrt{1+|x|^2}} \sigma_0(x).$$

$\square$

IV. L^1 CONVERGENCE

In this section, we shall show that our algorithm will yield an L^1 convergence for bounded domain, i.e. $\rho_R(x,\tau) = \lim_{k\to\infty} \rho_{k,R}(x,\tau)$, for $0 \le \tau \le T$, in the L^1 sense. For the convenience of the reader, we state the technique lemma will be used in the proof of Theorem 1.1 below.

Lemma 4.2: (Lemma 4.1, [9]) Let Ω be a bounded domain in $\mathbb{R}^n$ and let $v : \overline{\Omega} \times [0,T] \to \mathbb{R}$ be a C^1 function. Assume that $v(x,t) = 0$ for $(x,t) \in \partial\Omega \times [0,T]$. Let $\Omega_t^+ = \{x \in \Omega : v(x,t) \ge 0\}$. Then

$$\frac{d}{dt} \int_{\Omega_t^+} v(x,t) = \int_{\Omega_t^+} \frac{\partial v}{\partial t}(x,t),$$

for almost all $t \in [0,T]$.

Proof of Theorem 1.1: For the sake of brief notation, we omit the subscript Ω for ρ_Ω and $\rho_{i,\Omega}$ henceforth. Let $\Omega_t^+ = \{x \in \Omega : \rho(x,t) - \rho_i(x,t) \ge 0\}$. Apply Lemma 2.1 to $(\rho - \rho_i)$ taking place of ρ_R, with the test function $\psi \equiv 1$, we have

$$\frac{d}{dt} \int_{\Omega_t^+} (\rho - \rho_i) \le \int_{\Omega_t^+} (\rho - \rho_i) N + \int_{\Omega_t^+} \rho_i [N - N(\cdot;\tau_{i-1})], \tag{4.1}$$

by Lemma 4.2. All the boundary integrations, except $\int_{\Omega_t^+} D_w(\rho-\rho_i)\cdot\nu$, vanish, since $(\rho-\rho_i)|_{\partial\Omega_t^+}=0$. Moreover, $\int_{\Omega_t^+} D_w(\rho-\rho_i)\cdot\nu\le 0$ due to the similar argument in Proposition 3.2, where $\int_{\partial B_R} D_w\rho\cdot\nu\le 0$. Combine the conditions (1.16) and (1.17), (4.1) can be controlled by

$$\frac{d}{dt}\int_{\Omega_t^+}(\rho-\rho_i)\le C\int_{\Omega_t^+}(\rho-\rho_i)+\tilde{C}(t-\tau_{i-1})^\alpha\int_\Omega\rho. \tag{4.2}$$

To estimate $\int_\Omega\rho$, we apply Lemma 2.1 to ρ, with the test function $\psi\equiv 1$, we get

$$\frac{d}{dt}\int_\Omega\rho\le\int_\Omega\rho N\le C\int_\Omega\rho,$$

which implies

$$\int_\Omega\rho\le C\int_\Omega\sigma_{0,\Omega}, \tag{4.3}$$

where C is a generic constant, depending on T, for all $0\le t\le T$. Thus,

$$\frac{d}{dt}\int_{\Omega_t^+}(\rho-\rho_i)\le C\int_{\Omega_t^+}(\rho-\rho_i)+\tilde{C}(t-\tau_{i-1})^\alpha\int_\Omega\sigma_{0,\Omega}.$$

Multiply $e^{-\tilde{C}(t-\tau_{i-1})}$ on both sides and integrate from τ_{i-1} to t, we get

$$\int_{\Omega_t^+}(\rho-\rho_i)(x,t)\le e^{\tilde{C}(t-\tau_{i-1})}\int_{\Omega_{\tau_{i-1}}^+}(\rho-\rho_i)(x,\tau_{i-1})+C\frac{(t-\tau_{i-1})^{1+\alpha}}{1+\alpha}e^{\tilde{C}(t-\tau_{i-1})},$$

where C is a constant, which depends on T, $\int_\Omega\sigma_{0,\Omega}$. Similarly, one can also get that

$$\int_{\Omega_t^-}(\rho_i-\rho)(x,t)\le e^{\tilde{C}(t-\tau_{i-1})}\int_{\Omega_{\tau_{i-1}}^-}(\rho_i-\rho)(x,\tau_{i-1})+C\frac{(t-\tau_{i-1})^{1+\alpha}}{1+\alpha}e^{\tilde{C}(t-\tau_{i-1})}.$$

Consequently, we have

$$\begin{aligned}\int_\Omega|\rho-\rho_i|(x,t)\le&e^{\tilde{C}(t-\tau_{i-1})}\left[\int_\Omega|\rho-\rho_i|(x,\tau_{i-1})+C\frac{(t-\tau_{i-1})^{1+\alpha}}{1+\alpha}\right]\\\le&e^{\tilde{C}(t-\tau_{i-1})}\left[\int_\Omega|\rho-\rho_{i-1}|(x,\tau_{i-1})+C\frac{(t-\tau_{i-1})^{1+\alpha}}{1+\alpha}\right],\end{aligned} \tag{4.4}$$

since $\rho_i(x,\tau_{i-1})=\rho_{i-1}(x,\tau_{i-1})$, for $i=1,2,\cdots,k$. Applying (4.4) recursively, we obtain

$$\begin{aligned}\int_\Omega|\rho-\rho_k|(x,\tau_k)\le&e^{\tilde{C}(\tau_k-\tau_{k-1})}\left[\int_\Omega|\rho-\rho_{k-1}|(x,\tau_{k-1})+C\frac{(\tau_k-\tau_{k-1})^{1+\alpha}}{1+\alpha}\right]\\\le&e^{\tilde{C}T}\int_\Omega|\rho-\rho_0|(x,0)\\&+\frac{C}{1+\alpha}[(\tau_k-\tau_{k-1})^{1+\alpha}e^{\tilde{C}(\tau_k-\tau_{k-1})}+(\tau_{k-1}-\tau_{k-2})^{1+\alpha}e^{\tilde{C}(\tau_k-\tau_{k-2})}\\&+\cdots+(\tau_1-\tau_0)^{1+\alpha}e^{\tilde{C}(\tau_k-\tau_0)}]\\=&\frac{C}{1+\alpha}\frac{T^{1+\alpha}}{k^{1+\alpha}}\left(e^{\tilde{C}\frac{T}{k}}+e^{\tilde{C}\frac{2T}{k}}+\cdots+e^{\tilde{C}\frac{kT}{k}}\right)\le\frac{C}{k^\alpha},\end{aligned}$$

where C is a constant, which depends on α, T and $\int_\Omega \sigma_{0,\Omega}$. It is clear that $\int_\Omega |\rho - \rho_k| \to 0$, as $k \to \infty$. $\square$

V. LOWER BOUND ESTIMATE OF DENSITY FUNCTION

Theorem 5.5: Let ρ_R be the solution of (1.25), the robust DMZ equation on B_R. Assume that

1) $f(x,t)$ and $h(x,t)$ have at most polynomial growth in $|x|$, for all $t \in [0,T]$;
2) For any $0 \leq t \leq T$, there exists positive integer m and positive constants C' and C'' independent of R such that the following two inequalities hold on $\mathbb{R}^n$:

   (a) $$\frac{1}{2}\left[nm(m-2)\left|\left|GQG^T\right|\right|_\infty + m\,\mathrm{Tr}\left(GQG^T\right)\right]|x|^{m-2} - m|x|^{m-2}(f - D_wK)\cdot x + N(x,t) \geq -C'; \tag{5.1}$$

   (b) $$\left|n\left|\left|GQG^T\right|\right|_\infty\left(\frac{1}{2}m^2|x|^{2m-2} - m\left(\frac{1}{2}m - 1\right)|x|^{m-2}\right) - \frac{1}{2}m\,\mathrm{Tr}\left(GQG^T\right)|x|^{m-2} - m(f - D_wK)\cdot x|x|^{m-2}\right| \leq \frac{1}{2}nm(m+1)\left|\left|GQG^T\right|\right|_\infty |x|^{2m-2} + C'', \tag{5.2}$$

   where Tr$(*)$ is the trace of $*$.
3) Condition (1.16) is satisfied.

Then for any $R_0 < R$,

$$\begin{aligned}&\int_{B_{R_0}} \zeta\rho_R(x,T)\\ \geq &\frac{e^{(C-C')T - R_0^m}}{C'}\left(\frac{1}{2}nm(m+1)\left|\left|GQG^T\right|\right|_\infty R_0^{2m-2} + C''\right)\left(1 - e^{C'T}\right)\int_{B_R}\sigma_{0,R}(x)\\ &+ e^{-C'T}\int_{B_{R_0}}\zeta\sigma_{0,R}(x),\end{aligned}$$

where $\zeta(x) = e^{-\xi(x)} - e^{-\xi(R_0)}$, $\xi(x) = |x|^m$.

In particular, the solution ρ of the robust DMZ equation (1.5) on $\mathbb{R}^n$ has the estimate

$$\int_{\mathbb{R}^n} e^{-|x|^m}\rho(x,T) \geq e^{-C'T}\int_{\mathbb{R}^n} e^{-|x|^m}\sigma_0(x).$$

Proof: Apply Lemma 2.1 to ρ_R with the test function ψ to be $\zeta = e^{-\xi(x)} - e^{-\xi(R_0)}$, where $\xi(x)$ is an increasing function in $|x|$, we have

$$\frac{d}{dt}\int_{B_{R_0}}\zeta\rho_R = \int_{B_{R_0}}\rho_R\left[\frac{1}{2}D_w^2\zeta + (f - D_wK)\cdot\nabla\zeta + \zeta N\right].$$

All the boundary integrations vanish, since $\zeta|_{\partial B_R} = \rho_R|_{\partial B_R} = 0$. Direct computations yield that

$$\begin{aligned}
&\frac{d}{dt}\int_{B_{R_0}} \zeta\rho_R \\
=&\int_{B_{R_0}} \rho_R e^{-\xi(R_0)}\left[\frac{1}{2}\frac{\xi'^2(r)}{r^2}\sum_{i,j=1}^{n}\left(GQG^T\right)_{ij} x_i x_j - \frac{\xi'(r)}{r}(f - D_w K)\cdot x\right. \\
&\left.\qquad -\frac{1}{2}\sum_{i,j=1}^{n}\left(GQG^T\right)_{ij}\left[\left(\xi''(r) - \frac{\xi'(r)}{r}\right)\frac{x_i x_j}{r^2}\right] - \frac{1}{2}\mathrm{Tr}\left(GQG^T\right)\frac{\xi'(r)}{r}\right] \\
&+\int_{B_{R_0}} \zeta\rho_R\left[\frac{1}{2}D_w\xi\cdot\nabla\xi - \frac{1}{2}D_w^2\xi - (f - D_w K)\cdot\nabla\xi + N\right] \\
\triangleq&\,\mathrm{I}_5 + \int_{B_{R_0}} \zeta\rho_R[\mathrm{I}_6].
\end{aligned}$$

Let $\xi(r) = r^m$, where $r = |x|$, m is some positive integer sufficiently large. Through elementary computations, we get

$$\begin{aligned}
\mathrm{I}_6 =&\frac{1}{2}\frac{\xi'^2(r)}{r^2}\sum_{i,j=1}^{n}\left(GQG^T\right)_{ij} x_i x_j \\
&-\frac{1}{2}\left[m(m-2)r^{m-4}\sum_{i,j=1}^{n}\left(GQG^T\right)_{ij} x_i x_j + m r^{m-2}\mathrm{Tr}\left(GQG^T\right)\right] \\
&- m r^{m-2}(f - D_w K)\cdot x + N \\
\geq& -\frac{1}{2}\left[nm(m-2)\left|\left|GQG^T\right|\right|_\infty + m\,\mathrm{Tr}\left(GQG^T\right)\right] r^{m-2} - m r^{m-2}(f - D_w K)\cdot x + N \geq C',
\end{aligned}$$

where C' is a positive constant independent of R_0, by condition (5.1). For large enough m, we have

$$\begin{aligned}
|\mathrm{I}_5| \leq e^{-R_0^m}&\int_{B_R}\left| n\left|\left|GQG^T\right|\right|_\infty\left[\frac{1}{2}m^2 r^{2m-2} - m\left(\frac{1}{2}m - 1\right)r^{m-2}\right]\right. \\
&\qquad\left. -\frac{1}{2}m\,\mathrm{Tr}\left(GQG^T\right)r^{m-2} - m(f - D_w K)\cdot x r^{m-2}\right|\rho_R \\
\leq e^{-R_0^m}&\left(\frac{1}{2}nm(m+1)\left|\left|GQG^T\right|\right|_\infty R_0^{2m-2} + C''\right)\int_{B_R}\rho_R \\
\leq e^{CT-R_0^m}&\left(\frac{1}{2}nm(m+1)\left|\left|GQG^T\right|\right|_\infty R_0^{2m-2} + C''\right)\int_{B_R}\sigma_{0,R} \triangleq \gamma(R_0).
\end{aligned}$$

The last inequality follows by the similar argument of (4.3). Hence,

$$\frac{d}{dt}\int_{B_{R_0}}\zeta\rho_R \geq -\gamma(R_0) - C'\int_{B_{R_0}}\zeta\rho_R.$$

This implies

$$\int_{B_{R_0}} \zeta\rho_R(x,T) \geq e^{-C'T}\int_{B_{R_0}} \zeta\sigma_{0,R}(x) + \frac{\gamma(R_0)}{C'}\left(e^{-C'T}-1\right)$$

$$\geq e^{-C'T}\int_{B_{R_0}} \zeta\sigma_{0,R}(x)$$

$$+\frac{e^{(C-C')T-R_0^m}}{C'}\left(\frac{1}{2}nm(m+1)\left|\left|GQG^T\right|\right|_\infty R_0^{2m-2}+C''\right)\left(1-e^{C'T}\right)\int_{B_R}\sigma_{0,R}(x) \qquad (5.3)$$

Let $R_0 \to \infty$, we have

$$\int_{\mathbb{R}^n} e^{-|x|^m}\rho(x,T) \geq e^{-C'T}\int_{\mathbb{R}^n} e^{-|x|^m}\sigma_0(x).$$

$\square$

VI. NUMERICAL SIMULATIONS

We give the numerical simulations of the 1D cubic sensor:

$$\begin{cases} dx_t = dv_t \\ dy_t = x_t^3 dt + dw_t, \end{cases}$$

where x_t, $y_t \in \mathbb{R}$, v_t, w_t are scalar Brownian motion processes with $E[dv_t^T dv_t] = 1$, $E[dw_t^T dw_t] = 1$. The 1D Kolmogorov forward equation (1.13) here is

$$u_t = \frac{1}{2}u_{xx} - \frac{1}{2}x^6 u, \qquad (6.4)$$

for each time step. The updated initial data is

$$u_i(x,\tau_i) = e^{dy_t\frac{x^3}{4}}u_{i-1}(x,\tau_i).$$

We adopt the Hermite spectral method to get the approximate solution $\hat{u}_i$, for $i = 0, 1, \cdots, k$. $\hat{\rho}$ can be recovered by (1.12). The expectation of the state x_t is computed by definition

$$\mathbb{E}[x](t) = \frac{\int_{\mathbb{R}} x\hat{\rho}(x,t)dx}{\int_{\mathbb{R}} \hat{\rho}(x,t)dx}.$$

In Figure 1-2, we see that Yau-Yau's algorithm tracks the state's expectation very well, while the extended Kalman filter completely fails. Take the efficiency into consideration, it costs only around 7.5 seconds for Yau-Yau's algorithm to simulate $T = 10$, while the extended Kalman filter takes around 3 seconds in our PC. We are expecting some speedup if a more powerful computer is used. The potential advantage of Yau-Yau's algorithm is that multi-core processors can speedup



Fig. 1. Comparison of Yau-Yau's algorithm and extended Kalman filter for cubic sensor, with the initial condition $u_0(x) = e^{-\frac{x^4}{4}}$.



Fig. 2. Comparison of Yau-Yau's algorithm and extended Kalman filter for cubic sensor, with the initial condition $u_0(x) = e^{-\frac{x^4}{4}}(1 - \sin x)$.

further. For instance, if a dual-core processor is available, the computation of expectation of the state at the end of each time step can be handled by one processor, while the computation of the Fourier-Hermite coefficients is working on the other one. If more processors are provided, the first N Fourier-Hermite coefficients of the initial data at the beginning of each time step can be scattered to different processors and be computed simultaneously. In contrast with Yau-Yau's algorithm, the extended Kalman filter is only suitable programmed in sequential code, since the expectation and the covariance of the next time step are heavily relied on those of the previous time step. Even with multi-core processors, no speedup will be expected.

Since we directly deal with the density function, all the higher central moments of the state



Fig. 3. The 2nd to 6th central moments of the cubic sensor, with the initial condition $u_0(x) = e^{-\frac{x^4}{4}}$, computed by Yau-Yau's algorithm.

could be obtained by definition

$$\mathbb{E}[(x - \mathbb{E}[x])^k](t) = \frac{\int_{\mathbb{R}} (x - E[x_t])^k \hat{\rho}(x,t)dx}{\int_{\mathbb{R}} \hat{\rho}(x,t)dx},$$

for any $k \in \mathbb{Z}^+$. As we known, EKF is essentially obtained by truncating the first two moments to get the approximation. While in our algorithm, we don't truncate any order of moments. In Figure 3, we plot the 2^{nd} to 6^{th} central moments of the cubic sensor with the initial condition $u_0(x) = e^{-x^4/4}$.

APPENDIX

Before we show the existence of the weak solution, we'd like to give a priori estimations of up to the first order derivative of the solution to the robust DMZ equation on $B_R \times [0,T]$.

Theorem A.1: Consider the robust DMZ equation (1.25) on $\mathbb{Q}_R := B_R \times [0,T]$, where $B_R = \{x \in \mathbb{R}^n : |x| \leq R\}$ is a ball of radius R. Assume that

$$\left\| \frac{d}{dt}(GQG^T) \right\|_\infty < \infty, \tag{A.1}$$

for all $t \in [0,T]$. Suppose there exists a positive function $g(x)$ on $\mathbb{R}^n$ such that for all $t \in [0,T]$, g and $\tilde{g} \triangleq g + \log |D_w J|$ satisfies

1) $\left| D_w g + \frac{1}{2}\nabla(GQG^T) - F \right|^2 + 2\lambda_1 J \leq C,$ (A.2)

2) $D_w^2 g + 2D_w g \cdot \nabla g + 2[\nabla(GQG^T) - F] \cdot \nabla g + \frac{1}{2}\nabla^2(GQG^T) - \text{div}F + J \leq C,$ (A.3)

3) $D_w^2 \tilde{g} + 2D_w \tilde{g} \cdot \nabla \tilde{g} + 2[\nabla(GQG^T) - F] \cdot \nabla \tilde{g} + \frac{1}{2}\nabla^2(GQG^T) - \text{div}F + J \leq C,$ (A.4)

4) $\int_{\mathbb{R}^n} e^{2\tilde{g}}\sigma^2(x) \leq C$ and $\int_{\mathbb{R}^n} e^{2g} D_w\sigma \cdot \nabla\sigma \leq C,$ (A.5)

where C is a generic constant, which may differ from line to line, and $\nabla(*) = \left[\sum_{i=1}^n \frac{\partial(*)_{ij}}{\partial x_i}\right]_{j=1}^n$, $\nabla^2(*) = \sum_{i,j=1}^n \frac{\partial^2(*)_{ij}}{\partial x_i \partial x_j}$. Then, for $0 \leq t \leq T$,

$$\int_{B_R} e^{2g}\rho_R^2(x,t)dx \leq e^{Ct}\int_{B_R} e^{2g}\sigma^2(x)dx, \tag{A.6}$$

$$\int_{B_R} e^{2g} D_w\rho_R(x,t) \cdot \nabla\rho_R(x,t)dx \leq e^{Ct}\int_{B_R} e^{2g}D_w\sigma(x)\cdot\nabla\sigma(x)dx + Ce^{Ct}\int_{B_R} e^{2\tilde{g}}\sigma^2(x)dx, \tag{A.7}$$

where D_w and $J(x,t)$ is defined in (2.4)and (1.9), respectively.

Remark A.1: The conditions in Theorem A.1 are easily checked, if the drift terms $h(x)$ and $f(x)$ are at most polynomial growth in $r = |x|$. However, in general, the existence of such g is not always available.

Proof: Let g be some positive function on $\mathbb{R}^n$.

$$\begin{aligned}\frac{d}{dt}\int_{B_R} e^{2g}\rho_R^2 =& \int_{B_R} e^{2g}\rho_R D_w^2\rho_R + 2\int_{B_R} e^{2g}\rho_R(F\cdot\rho_R) + 2\int_{B_R} e^{2g}J\rho_R^2 \\ \triangleq& \mathrm{I} + \mathrm{II} + \mathrm{III}.\end{aligned} \tag{A.8}$$

Apply integration by parts to I and II in (A.8)

$$\begin{aligned}\mathrm{I} =& -2\int_{B_R}\rho_R e^{2g}D_w g\cdot\nabla\rho_R - \int_{B_R} e^{2g}D_w\rho_R\cdot\nabla\rho_R - \int_{B_R} e^{2g}\rho_R\nabla(GQG^T)\cdot\nabla\rho_R \\ \leq& -2\int_{B_R}\rho_R e^{2g}D_w g\cdot\nabla\rho_R - \int_{B_R} e^{2g}\rho_R\nabla(GQG^T)\cdot\nabla\rho_R \triangleq \mathrm{I}_1 + \mathrm{I}_2.\end{aligned}$$

Integration by parts further, we have

$$\begin{aligned}\mathrm{I}_1 =& 4\int_{B_R} e^{2g}\rho_R^2 D_w g\cdot\nabla g + 2\int_{B_R} e^{2g}\rho_R D_w g\cdot\nabla\rho_R + 2\int_{B_R} e^{2g}\rho_R^2\nabla(GQG^T)\cdot\nabla g \\ &+ 2\int_{B_R} e^{2g}\rho_R^2 D_w^2 g.\end{aligned} \tag{A.9}$$

Notice that the second term of the right-hand side of (A.9) is $-\mathrm{I}_1$, we have

$$\mathrm{I}_1 = 2\int_{B_R} e^{2g}\rho_R^2 D_w g\cdot\nabla g + \int_{B_R} e^{2g}\rho_R^2[\nabla(GQG^T)\cdot\nabla g + D_w^2 g]. \tag{A.10}$$

The similar argument applies to I_2:

$$\mathrm{I}_2 = \int_{B_R} e^{2g}\rho_R^2 \nabla(GQG^T)\cdot\nabla g + \frac{1}{2}\int_{B_R} e^{2g}\rho_R^2\nabla^2(GQG^T). \tag{A.11}$$

Thus,

$$\mathrm{I} \leq \int_{B_R} e^{2g}\rho_R^2\left[D_w^2 g + 2D_w g\cdot\nabla g + 2\nabla(GQG^T)\cdot\nabla g + \frac{1}{2}\nabla^2(GQG^T)\right]. \tag{A.12}$$

The same trick of I_1 applies to II in (A.8), we obtain

$$\mathrm{II} = -\int_{B_R} e^{2g}\rho_R^2[2F\cdot\nabla g + \mathrm{div}F]. \tag{A.13}$$

Substitute (A.12) and (A.13) back to (A.8), we obtain

$$\begin{aligned}
&\frac{d}{dt}\int_{B_R} e^{2g}\rho_R^2\\
\leq&\int_{B_R} e^{2g}\rho_R^2\left\{D_w^2 g + 2D_w g\cdot\nabla g + 2[\nabla(GQG^T) - F]\cdot\nabla g + \frac{1}{2}\nabla^2(GQG^T) - \mathrm{div}F + J\right\}\\
\leq&C\int_{B_R} e^{2g}\rho_R^2,
\end{aligned}$$

by condition (A.3). (A.6) follows directly from Gronwall's inequality. To show (A.7), we consider

$$\begin{aligned}
&\frac{d}{dt}\int_{B_R} e^{2g}D_w\rho_R\cdot\nabla\rho_R\\
=&\int_{B_R} e^{2g}\sum_{i,j=1}^{n}\frac{d}{dt}(GQG^T)_{ij}\frac{\partial\rho_R}{\partial x_i}\frac{\partial\rho_R}{\partial x_j} + 2\int_{B_R} e^{2g}\sum_{i,j=1}^{n}(GQG^T)_{ij}\frac{\partial}{\partial x_i}\left(\frac{\partial\rho_R}{\partial t}\right)\frac{\partial\rho_R}{\partial x_j}\\
\triangleq&\mathrm{IV} + \mathrm{V}.
\end{aligned} \tag{A.14}$$

Due to condition (A.1), IV of (A.14) turns out to be

$$\begin{aligned}
\mathrm{IV} \leq&\frac{1}{2}\left\|\frac{d}{dt}(GQG^T)\right\|_\infty\int_{B_R} e^{2g}\sum_{i,j=1}^{n}\left[\left(\frac{\partial\rho_R}{\partial x_i}\right)^2 + \left(\frac{\partial\rho_R}{\partial x_j}\right)^2\right] = n\left\|\frac{d}{dt}(GQG^T)\right\|_\infty\int_{B_R} e^{2g}|\nabla\rho_R|^2\\
\leq&\frac{n}{\lambda_1}\left\|\frac{d}{dt}(GQG^T)\right\|_\infty\int_{B_R} e^{2g}D_w\rho_R\cdot\nabla\rho_R,
\end{aligned} \tag{A.15}$$

since $D_w\rho_R \cdot \nabla\rho_R \geq \lambda_1|\nabla\rho_R|^2$. Next, V in (A.14) is

$$\begin{aligned}
\mathrm{V} =& -2\int_{B_R} e^{2g}[(2D_w g + \nabla(GQG^T))\cdot\nabla\rho_R + D_w^2\rho_R]\cdot\left(\frac{1}{2}D_w^2\rho_R + F\cdot\nabla\rho_R + J\rho_R\right)\\
=& -\int_{B_R} e^{2g}\left\{D_w^2\rho_R + \left[D_w g + \frac{1}{2}\nabla(GQG^T) + F\right]\cdot\nabla\rho_R\right\}^2\\
&+\int_{B_R} e^{2g}\left[D_w g + \frac{1}{2}\nabla(GQG^T) - F\right]^2|\nabla\rho_R|^2\\
&-2\int_{B_R} e^{2g}[D_w^2\rho_R + (2D_w g + \nabla(GQG^T))\cdot\nabla\rho_R]J\rho_R\\
\leq& \int_{B_R} e^{2g}\left[D_w g + \frac{1}{2}\nabla(GQG^T) - F\right]^2|\nabla\rho_R|^2\\
&-2\int_{B_R} e^{2g}[D_w^2\rho_R + (2D_w g + \nabla(GQG^T))\cdot\nabla\rho_R]J\rho_R. \qquad (A.16)
\end{aligned}$$

Notice that

$$\begin{aligned}
\int_{B_R} e^{2g}D_w^2\rho_R J\rho_R = -\int_{B_R} e^{2g}\big[&2(D_w g\cdot\nabla\rho_R)J\rho_R + JD_w\rho_R\cdot\nabla\rho_R\\
&+(D_w\rho_R\cdot\nabla J)\rho_R + \nabla(GQG^T)\cdot\nabla\rho_R J\rho_R\big]. \qquad (A.17)
\end{aligned}$$

Take (A.17) into account, V becomes

$$\mathrm{V} \leq \int_{B_R} e^{2g}\left\{\frac{1}{\lambda_1}\left\{\left[D_w g + \frac{1}{2}\nabla(GQG^T) - F\right]^2 + 1\right\} + 2J\right\}D_w\rho_R\cdot\nabla\rho_R + \int_{B_R} e^{2g}|D_w J|^2\rho_R^2. \qquad (A.18)$$

Combine (A.15) and (A.18), we have

$$\begin{aligned}
&\frac{d}{dt}\int_{B_R} e^{2g}D_w\rho_R\cdot\nabla\rho_R\\
&\leq \int_{B_R} e^{2g}\left\{\frac{1}{\lambda_1}\left\{n\left\|\frac{d}{dt}(GQG^T)\right\|_\infty + \left[D_w g + \frac{1}{2}\nabla(GQG^T) - F\right]^2 + 1\right\} + 2J\right\}D_w\rho_R\cdot\nabla\rho_R\\
&\quad + \int_{B_R} e^{2g}|D_w J|^2\rho_R^2. \qquad (A.19)
\end{aligned}$$

By conditions (A.2)-(A.5), the estimate (A.7) follows immediately. ☐

Proof of Theorem 2.3: Let R_k be a sequence of positive number such that $\lim_{k\to\infty} R_k = \infty$. Let $\rho_k(x,t)$ be the solution of the robust DMZ equation (1.25) on $B_{R_k}\times[0,T]$, where $B_{R_k} = \{x\in\mathbb{R}^n : |x| \leq R_k\}$ is a ball of radius R_k. In view of Theorem A.1, the sequence $\{\rho_k\}$ is a bounded set in $H_0^{1;1}(\mathbb{Q}_{R_k})$. Thus, there exists a subsequence $\{\rho_{k'}\}$ which is weakly convergent to ρ. Moreover, ρ has the weak derivative $\frac{\partial \rho}{\partial x_i} \in L^2(\mathbb{Q}_{R_k})$, and $\frac{\partial \rho_{k'}}{\partial x_i}$ weakly tends to it. Now we claim that the weak derivative $\frac{\partial \rho}{\partial t}$ exists. To see this, let $\Phi(x,t) \in H_0^{1;1}(\mathbb{Q}_{R_k})$, then

$$
\begin{aligned}
&\iint_{\mathbb{Q}_{R_k}} \frac{1}{2} \sum_{i,j=1}^{n} (GQG^T)_{ij} \frac{\partial \Phi}{\partial x_j} \frac{\partial \rho}{\partial x_i} + \left[\sum_{i=1}^{n} \left(\sum_{j=1}^{n} \frac{\partial (GQG^T)_{ij}}{\partial x_j} - F_i \right) \frac{\partial \rho}{\partial x_i} - J\rho \right] \Phi \\
&\qquad = \lim_{k' \to \infty} \iint_{\mathbb{Q}_{R_k}} \frac{1}{2} \sum_{i,j=1}^{n} (GQG^T)_{ij} \frac{\partial \Phi}{\partial x_j} \frac{\partial \rho_{k'}}{\partial x_i} + \left[\sum_{i=1}^{n} \left(\sum_{j=1}^{n} \frac{\partial (GQG^T)_{ij}}{\partial x_j} - F_i \right) \frac{\partial \rho_{k'}}{\partial x_i} - J\rho_{k'} \right] \Phi \\
&\qquad = - \lim_{k' \to \infty} \iint_{\mathbb{Q}_{R_k}} \frac{\partial \rho_{k'}}{\partial t} \Phi = \lim_{k' \to \infty} \iint_{\mathbb{Q}_{R_k}} \rho_{k'} \frac{\partial \Phi}{\partial t} = \iint_{\mathbb{Q}_{R_k}} \rho \frac{\partial \Phi}{\partial t}.
\end{aligned}
$$

Clearly, $\rho(x,0) = \lim_{k' \to \infty} \rho_{k'}(x,0) = \sigma_0(x)$.

To show the uniqueness of the solution, we only need to show that $\rho(x,t) = 0$ on $\mathbb{Q}_T$ if $\rho(x,0) = 0$. Let $\alpha T < 1$. For any test function $\psi(x,t) = e^{cr}\Phi(x,t)$, where $r = |x|$, c is some constant and $\Phi(x,t) \in H_0^{1;1}(\mathbb{Q}_T)$, then $\rho(x,t)$ satisfies

$$
\begin{aligned}
&\int_{\mathbb{R}^n} \rho(x,T)\Phi(x,T)e^{cr}dx - \int_0^T \int_{\mathbb{R}^n} \rho(x,t) \frac{\partial \Phi}{\partial t}(x,t) e^{cr} dx dt \\
= &\int_{\mathbb{Q}_T} -\frac{1}{2} e^{cr} \nabla \Phi(x,t) \cdot D_w \rho(x,t) - \frac{c}{2} e^{cr} \Phi(x,t) \nabla r \cdot D_w \rho(x,t) + \tilde{F}(x,t) \cdot \nabla \rho(x,t) \Phi(x,t) e^{cr} \\
&\qquad + J(x,t)\rho(x,t)\Phi(x,t)e^{cr} dx dt.
\end{aligned}
\tag{A.20}
$$

where $\tilde{F}$ is defined in (2.2). Approximate $\rho(x,t)$ by $\Phi(x,t)$ in the $H^{1;1}(\mathbb{Q}_T)$-norm, we get

$$
\begin{aligned}
&\int_{\mathbb{R}^n} \rho^2(x,T) e^{cr} dx \\
= &\int_{\mathbb{Q}_T} e^{cr} \left[-D_w \rho(x,t) \cdot \nabla \rho(x,t) - c\rho(x,t) \nabla r \cdot D_w \rho(x,t) \right. \\
&\qquad \left. + (\tilde{F}(x,t) + F(x,t)) \cdot \nabla \rho(x,t) \rho(x,t) + 2J(x,t)\rho^2(x,t) \right] dx dt. \\
\leq &\int_{\mathbb{Q}_T} e^{cr} [-\lambda_1 |\nabla \rho(x,t)|^2 - c\rho(x,t) D_w r \cdot \nabla \rho(x,t) \\
&\qquad + (F(x,t) + \tilde{F}(x,t)) \cdot \nabla \rho(x,t) \rho(x,t) + 2J(x,t)\rho^2(x,t)] dx dt. \\
= &- \lambda_1 \int_{\mathbb{Q}_T} e^{cr} \left\{ |\nabla \rho(x,t)| + \frac{1}{2\lambda_1} [cD_w r - (F(x,t) + \tilde{F}(x,t))] \rho(x,t) \right\}^2 dx dt \\
&+ \int_{\mathbb{Q}_T} e^{cr} \left\{ 2J(x,t) - \frac{1}{4\lambda_1} [cD_w r - (F(x,t) + \tilde{F}(x,t))]^2 \right\} \rho^2(x,t) dx dt \\
\leq &\int_{\mathbb{Q}_T} e^{cr} \left\{ 2J(x,t) - \frac{1}{4\lambda_1} [cD_w r - (F(x,t) + \tilde{F}(x,t))]^2 \right\} \rho^2(x,t) dx dt,
\end{aligned}
\tag{A.21}
$$

due to the positive definite of (GQG^T). By condition (2.1), we have

$$\int_{\mathbb{R}^n} e^{cr}\rho^2(x,T)dx \leq \alpha \int_{\mathbb{Q}_T} e^{cr}\rho^2(x,t)dxdt. \tag{A.22}$$

According to the mean value theorem, there exists $T_1 \in (0,T)$ such that

$$\int_{\mathbb{Q}_T} e^{cr}\rho^2(x,t)dxdt = \int_0^T \int_{\mathbb{R}^n} e^{cr}\rho^2(x,t)dxdt = T\int_{\mathbb{R}^n} e^{cr}\rho^2(x,T_1)dx. \tag{A.23}$$

Apply (A.22) and (A.23) recursively, there exists $T_m \in (0,T)$ such that

$$\int_{\mathbb{R}^n} e^{cr}\rho^2(x,T)dx \leq (\alpha T)^m \int_{\mathbb{R}^n} e^{cr}\rho^2(x,T_m)dx.$$

Since $\alpha T < 1$, we conclude that $\rho(x,t) \equiv 0$ for a.e $(x,t) \in \mathbb{Q}_T$. $\square$

REFERENCES

[1] A. Bensoussan, R. Glowinski and A. Rascanu, *Approximation of the Zakai equation by the splitting up method*, SIAM J. Control Optim., **28**(1990), pp 1420-1431.

[2] T. E. Duncan, *Probability densities for diffusion processes with applications to nonlinear filtering theory*, Ph.D. thesis, Stanford, 1967.

[3] W. Fleming and S. Mitter, *Optimal control and nonlinear filtering for nondegenerate diffusion processes*, Stochastics, **8**(1982), pp 63-77.

[4] A. Friedman, *Partial differential equations of parabolic type*, Prentice-Hall, Englewood Cliffs, NJ, 1964.

[5] R. E. Mortensen, *Optimal control of continuous time stochastic systems*, Ph.D. thesis, University of California, Berkeley, CA, USA, 1996.

[6] E. Pardoux, *Stochastic partial differential equations and filtering of diffusion processes*, Stochastics, **3**(1979), pp 127-167.

[7] B. L. Rozovsky, *Stochastic partial differential equations arising in nonlinear filtering problems*, Usp. Mat. Nauk., **27**(1972), pp. 213-214.

[8] S. L. Sobolev, *Applications of functional analysis in mathematical physics*, Tran. Math. Monographs 7, AMS, Providence, RI, 1963.

[9] S. S.-T. Yau, S.-T. Yau, *Real time solution of nonlinear filtering problem without memory II*, SIAM Journal of Control and Optimization, **47**:1(2008), pp. 163-195.

[10] M. Zakai, *On the optimal filtering of diffusion processes*, Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete, **11**(1969), pp. 230-243.